
CLIMATIC ATLAS OF ICING POTENTIAL OVER NORTH AMERICA
(U) AIR FORCE ENVIRONMENTAL TECHNICAL APPLICATIONS
CENTER SCOTT AFB IL JAN 86 USAFETAC/DS-86/881 AD-A174 268 1/2 UNCLASSIFIED F/G 4/1 NL



CROCOPY RESOLUTION TEST CHART
MATIONAL BUREAU OF STANDARDS-1963-A





USAFETAC/DS-86/001

CLIMATIC ATLAS OF ICING POTENTIAL OVER NORTH AMERICA

JANUARY 1986



Approved For Public Release; Distribution Unlimited

UNITED STATES AIR FORCE AIR WEATHER SERVICE (MAC) USAF ENVIRONMENTAL TECHNICAL APPLICATIONS CENTER

SCOTT AIR FORCE BASE, ILLINOIS 62225-5458

86 11 18 087

REPORT DOCUMENTATION PAGE

- 1a. Report Security Classification: UNCLASSIFIED
- 3. <u>Distribution/Availability of Report:</u> Approved for public release; distribution is unlimited.
- 4. Performing Organization Report Number: USAFETAC/DS-86/001
- 6a. Name of Performing Organization: USAFTIAC
- 6b. Office Symbol: ECO
- 6c. Address: Scott AFB, IL 62225-5438
- 11. Title: A Climatic Atlas of Icing Potential Over North America (UNCLASSIFIED)
- 13a. Type of Report: Data Summary
- 14. Date of Report: January 1936
- 15. Page Count: 126
- 17. COSATI Codes: Field--04, Group--02
- 18. Subject Terms: *CLIMATE, *CLIMATOLOGY, *AIRCRAFT ICING, *ATLAS, *NORTH AMERICA, SMITH-FEDDES, *METEOROLOGY, satellite meteorology, liquid water content, 3DNEPH.
- Abstract: A climatic atlas of charts that show percent frequency of occurrence of potential loing conditions over North America. Since loing reports are sparse, an accurate loing climatology is difficult to obtain. A climatology of elements for potential loing conditions, however (such as liquid water content and temperature), can be derived. The Smith-Feddes Liquid Water Content (LWC) computer model uses combined data from the Air Force Global Weather Central (AFGWC); Three-Dimensional Nephanalysis Data Base (3DNEPH) and the upper air Analysis Data Set (ADS) for input. Output consists of percent frequency of occurrence of a liquid water content threshold concentration in a defined layer and temperature range. Output was generated for three concentrations, three layers and for temperatures at or below freezing. This data was then manually converted into 117 monthly and annual maps showing percent frequency of occurrence for potential loing conditions.
- 20. Distribution/Availability of Abstract: Same as report.
- 21. Abstract Security Classification: UNCLASSIFIED
- 22a. Name of Responsible Individual: 1 Lt Paul R. Thorson
- 22b. Telephone: 618 256~2642
- 22c. Office Symbol: ECO

4

	Accession For
	NTES GRAZI
	DT.C TAB
-	Unampurced /
	July 100% Title -
Ą	
	By
	Distribution/
į	Availability Codes
	ps 11 and/or
	Dist Decial
-	
	\mathcal{U}

DD FORM 1473

1,000 pt 1,0

PREFACE

This document was completed for USAFETAC Project 3702, initiated to satisfy a 5th Weather Wing request for icing climatology. SMSgt Charles Travers, USAFETAC/ECO, started the project in April 1985 and was the project manager until his transfer to another branch in August 1985. He was succeeded as project manager by 1st Lt Paul Thorson, who saw the project through to its completion. All manual chart contouring was by SSgt Charles Carrier. Capt James Millard was the technical consultant. TSgt Ronald Coleman wrote computer software that simplified data extraction from the liquid water content database. The USAFETAC Data Automation Branch also deserves recognition for dealing with the considerable computer time requirements for the Smith-Feddes liquid water content model.





CONTENTS



PURPOSE AND BACKGROUND										
THE SMITH-FEDDES LWC MODEL										
How the	LWC Model W	orks1								
PERIOD OF RECORD										
CHART PREPARATION METHOD										
USING THE CHARTS2										
EXAMPLE CHA	ART INTERPRE	TATION2								
CHART 1	JANUARY	Surface to 1,524 Meters, Concentration $.10 \text{ g/m}^3$								
CHART 2	FEBRUARY	Surface to 1,524 Meters, Concentration $.10 \text{ g/m}^34$								
CHART 3	MARCH	Surface to 1,524 Meters, Concentration $.10 \text{ g/m}^3$								
CHART 4	APRIL	Surface to 1,524 Meters, Concentration $.10 \text{ g/m}^3$ 6								
CHART 5	MAY	Surface to 1,524 Meters, Concentration $.10 \text{ g/m}^3$ 7								
CHART 6	JUNE	Surface to 1,524 Meters, Concentration $.10 \text{ g/m}^38$								
CHART 7	JULY	Surface to 1,524 Meters, Concentration .10 g/m ³ 9								
CHART 8	AUGUST	Surface to 1,524 Meters, Concentration $.10 \text{ g/m}^310$								
CHART 9	SEPTEMBER	Surface to 1,524 Meters, Concentration .10 g/m ³ 11								
CHART 10	OCTOBER	Surface to 1,524 Meters, Concentration $.10 \text{ g/m}^3$								
CHART 11	NOVEMBER	Surface to 1,524 Meters, Concentration .10 g/m ³ 13								
CHART 12	DECEMBER	Surface to 1,524 Meters, Concentration .10 g/m ³ 14								
CHART 13	ANNUAL	Surface to 1,524 Meters, Concentration .10 g/m ³ 15								
CHART 14	JANUARY	Surface to 1,524 Meters, Concentration $.50 \text{ g/m}^3$								
CHART 15	FEBRUARY	Surface to 1,524 Meters, Concentration .50 g/m ³ 17								
CHART 16	MARCH	Surface to 1,524 Meters, Concentration .50 g/m ³ 18								
CHART 17	APRIL	Surface to 1,524 Meters, Concentration .50 g/m ³ 19								
CHART 18	MAY	Surface to 1,524 Meters, Concentration $.50 \text{ g/m}^3$ 20								
CHART 19	JUNE	Surface to 1,524 Meters, Concentration $.50 \text{ g/m}^3$ 21								
CHART 20	JULY	Surface to 1,524 Meters, Concentration .50 g/m ³ 22								
CHART 21	AUGUST	Surface to 1,524 Meters, Concentration $.50 \text{ g/m}^3$ 23								
CHART 22	SEPTEMBER	Surface to 1,524 Meters, Concentration .50 g/m ³ 24								
CHART 23	OCTOBER	Surface to 1,524 Meters, Concentration .50 g/m ³ 25								
CHART 24	NOVEMBER	Surface to 1,524 Meters, Concentration .50 g/m ³ 26								
CHART 25	DECEMBER	Surface to 1,524 Meters, Concentration .50 g/m ³ 27								
CHART 26	ANNUAL	Surface to 1,524 Meters, Concentration .50 g/m ³ 28								



CHART 27	JANUARY	Surface to 1,524 Meters, Concentration 1.00 g/m ³ 29
CHART 28	FEBRUARY	Surface to 1,524 Meters, Concentration 1.00 g/m^3 30
CHART 29	MARCH	Surface to 1,524 Meters, Concentration 1.00 g/m ³ 39
CHART 30	APRIL	Surface to 1,524 Meters, Concentration 1.00 g/m ³
CHART 31	MAY	Surface to 1,524 Meters, Concentration 1.00 g/m^3
CHART 32	JUNE	Surface to 1,524 Meters, Concentration 1.00 g/m ³
CHART 33	JULY	Surface to 1,524 Meters, Concentration 1.00 g/m^3 35
CHART 34	AUGUST	Surface to 1,524 Meters, Concentration 1.00 g/m^3
CHART 35	SEPTEMBER	Surface to 1,524 Meters, Concentration 1.00 g/m ³ 37
CHART 36	OCTOBER	Surface to 1,524 Meters, Concentration 1.00 g/m ³
CHART 37	NOVEMBER	Surface to 1,524 Meters, Concentration 1.00 g/m ³ 39
CHART 38	DECEMBER	Surface to 1,524 Meters, Concentration 1.00 g/m ³ 40
CHART 39	ANNUAL	Surface to 1,524 Meters, Concentration 1.00 g/m ³ 41
CHART 40	JANUARY	1,525 to 3,048 Meters, Concentration .10 g/m ³
CHART 41	FEBRUARY	1,525 to 3,048 Meters, Concentration .10 g/m ³ 43
CHART 42	MARCH	1,525 to 3,048 Meters, Concentration .10 g/m^3 41
CHART 43	APRIL	1,525 to 3,048 Meters, Concentration .10 g/m ³ 45
CHART 44	MAY	1,525 to 3,048 Meters, Concentration .10 g/m^3 46
CHART 45	JUNE	1,525 to 3,048 Meters, Concentration .10 g/m^3 47
CHART 46	JULY	1,525 to 3,048 Meters, Concentration .10 g/m^3 48
CHART 47	AUGUST	1,525 to 3,048 Meters, Concentration .10 g/m ³ 49
CHART 48	SEPTEMBER	1,525 to 3,048 Meters, Concentration .10 g/m ³
CHART 49	OCTOBER	1,525 to 3,048 Meters, Concentration .10 g/m ³ 51
CHART 50	NOVEMBER	1,525 to 3,048 Meters, Concentration .10 g/m^3 52
CHART 51	DECEMBER	1,525 to 3,048 Meters, Concentration $.10 \text{ g/m}^3$
CHART 52	ANNUAL	1,525 to 3,048 Meters, Concentration .10 g/m ³ 51
CHART 53	JANUARY	1,525 to 3,048 Meters, Concentration .50 g/m^3 55
CHART 54	FEBRUARY	1,525 to 3,048 Meters, Concentration .50 g/m^3
CHART 55	MARCH	1,525 to 3,048 Meters, Concentration .50 g/m^3 51
CHART 56	APRIL	1,525 to 3,048 Meters, Concentration .50 g/m^3 58
CHART 57	MAY	1,525 to 3,048 Meters, Concentration .50 g/m 3 59
CHART 58	JUNE	1,525 to 3,048 Meters, Concentration .50 g/m 3 60
CHART 59	JULY	1,525 to 3,048 Meters, Concentration .50 g/m^3 61
CHART 60	AUGUST	1,525 to 3,048 Meters, Concentration .50 g/m^3
CHAPT 61	CROTEMBED	1.525 to 2.088 Metans, Concentration, 50 g/m ³





CHART 62	OCTOBER	1,525 to 3,048 Meters, Concentration .50 g/m ³ 64
CHART 63	NOVEMBER	1,525 to 3,048 Meters, Concentration .50 g/m ³ 65
CHART 64	DECEMBER	1,525 to 3,048 Meters, Concentration .50 g/m ³
CHART 65	ANNUAL	1,525 to 3,048 Meters, Concentration .50 g/m ³ 67
CHART 66	JANUARY	1,525 to 3,048 Meters, Concentration 1.00 g/m ³
CHART 67	FEBRUARY	1,525 to 3,048 Meters, Concentration 1.00 g/m ³ 69
CHART 68	MARCH	1,525 to 3,048 Meters, Concentration 1.00 g/m ³ 70
CHART 69	APRIL	1,525 to 3,048 Meters, Concentration 1.00 g/m ³ 71
CHART 70	MAY	1,525 to 3,048 Meters, Concentration 1.00 g/m ³ 72
CHART 71	JUNE	1,525 to 3,048 Meters, Concentration 1.00 g/m ³
CHART 72	JULY	1,525 to 3,048 Meters, Concentration 1.00 g/m ³
CHART 73	AUGUST	1,525 to 3,048 Meters, Concentration 1.00 g/m ³
CHART 74	SEPTEMBER	1,525 to 3,048 Meters, Concentration 1.00 g/m ³ 76
CHART 75	OCTOBER	1,525 to 3,048 Meters, Concentration 1.00 g/m^3 77
CHART 76	NOVEMBER	1,525 to 3,048 Meters, Concentration 1.00 g/m ³ 78
CHART 77	DECEMBER	1,525 to 3,048 Meters, Concentration 1.00 g/m^3 79
CHART 78	ANNUAL	1,525 to 3,048 Meters, Concentration 1.00 g/m^3 80
CHART 79	JANUARY	3,049 to 4,572 Meters, Concentration $.10 \text{ g/m}^381$
CHART 80	FEBRUARY	3,049 to 4,572 Meters, Concentration $.10 \text{ g/m}^382$
CHART 81	MARCH	3,049 to 4,572 Meters, Concentration $.10 \text{ g/m}^3$ 83
CHART 82	APRIL	3,049 to 4,572 Meters, Corcentration .10 g/m^3 84
CHART 83	MAY	3,049 to 4,572 Meters, Concentration $.10 \text{ g/m}^385$
CHART 84	JUNE	3,049 to 4,572 Meters, Concentration $.10 \text{ g/m}^3$ 86
CHART 85	JULY	3,049 to 4,572 Meters, Concentration $.10 \text{ g/m}^387$
CHART 86	AUGUST	3,049 to 4,572 Meters, Concentration $.10 \text{ g/m}^3$ 88
CHART 87	SEPTEMBER	3,049 to 4,572 Meters, Concentration $.10 \text{ g/m}^389$
CHART 88	OCTOBER	3,049 to 4,572 Meters, Concentration $.10 \text{ g/m}^390$
CHART 89	NOVEMBER	3,049 to 4,572 Meters, Concentration $.10 \text{ g/m}^3$
CHART 90	DECEMBER	3,049 to 4,572 Meters, Concentration .10 g/m^3 92
CHART 91	ANNUAL	3,049 to 4,572 Meters, Concentration $.10 \text{ g/m}^393$
CHART 92	JANUARY	3,049 to 4,572 Meters, Concentration .50 g/m^3 94
CHART 93	FEBRUARY	3,049 to 4,572 Meters, Concentration .50 g/m^3 95
CHART 94	MARCH	3,049 to 4,572 Meters, Concentration .50 g/m^3 96
CHART 95	APRIL	3,049 to 4,572 Meters, Concentration .50 g/m^3 97
CHART 96	MAY	3,049 to 4,572 Meters, Concentration .50 g/m^3 98



CHART	97	JUNE	3,049 to 4,572 Meters, Concentr tion .50 g/m ³ 99	
CHART	98	JULY	3,049 to 4,572 Meters, Concentration .50 g/m^3	
CHART	99	AUGUST	3,049 to 4,572 Meters, Concentration .50 g/m^3	
CHART	100	SEPTEMBER	3,049 to 4,572 Meters, Concentration .50 g/m^3	
CHART	101	OCTOBER	3,049 to 4,572 Meters, Concentration .50 g/m^3	
CHART	102	NOVEMBER	3,049 to 4,572 Meters, Concentration .50 g/m^3	
CHART	103	DECEMBER	3,049 to 4,572 Meters, Concentration .50 g/m 3	
CHART	104	ANNUAL	3,049 to 4,572 Meters, Concentration .50 g/m^3	
CHART	105	JANUARY	3,049 to 4,572 Meters, Concentration 1.00 g/m 3	
CHART	106	FEBRUARY	3,049 to 4,572 Meters, Concentration 1.00 g/m 3	
CHART	107	MARCH	3,049 to 4,572 Meters, Concentration 1.00 g/m 3	
CHART	108	APRIL	3,049 to 4,572 Meters, Concentration 1.00 g/m 3 110	
CHART	109	MAY	3,049 to 4,572 Meters, Concentration 1.00 g/m 3 111	
CHART	110	JUNE	3,049 to 4,572 Meters, Concentration 1.00 g/m^3	
CHART	111	JULY	3,049 to 4,572 Meters, Concentration 1.90 g/m 3 113	
CHART	112	AUGUST	3,049 to 4,572 Meters, Concentration 1.00 g/m^3	
CHART	113	SEPTEMBER	3,049 to 4,572 Meters, Concentration 1.00 g/m 3	
CHART	114	OCTOBER	3,049 to 4,572 Meters, Concentration 1.00 g/m^3	
CHART	115	NOVEMBER	3,049 to 4,572 Meters, Concentration 1.00 g/m^3	
CHART	116	DECEMBER	3,049 to 4,572 Meters, Concentration 1.00 g/m^3	
CHART	117	ANNUAL	3,049 to 4,572 Meters, Concentration 1.00 g/m^3 119	
BIBLIOGRAPHY120				
DISTRIBUTION 121				



PURPOSE AND BACKGROUND

The Requirement. This atlas is based on a 5th Weather Wing request for North American aircraft icing climatology to use in close air support and other low-level operations. Unfortunately, the science of forecasting icing is far from exact and reports of aircraft icing were scarce. In addition, aircraft icing is a function of airframe geometry, airspeed, flight profile, and degree of exposure-none of which can be quantified or averaged to produce a meaningful icing climatology. For all these reasons, production of an accurate icing climatology was a challenge. Nonetheless, the requirement was genine: there was a definite need for a simple and universally applicable climatology related to aircraft icing.

The Solution. USAFFTAC/ESS chose to fulfill the requirement with preparation of this atlas, which provides graphic presentation of aircraft icing potential, given certain specified conditions. The four elements that determine potential icing conditions are (1) liquid water content, (2) cloud drop-size distribution, (3) cloud temperature, and (4) ambient temperature. Of these, liquid water content and ambient temperature can be derived and quantified. The tool used to produce the charts in this atlas was the Smith-Feddes Liquid Water Content (LWC) model, which uses the Air Force Global Weather Central (AFGWC) Three-Dimensional Nephanalysis Data Base (3DNEPH) and the upper air Analysis Data Set (ADS) for input. The combined data sets, using satellite meteorological data, surface observations, and rawinsonde observations, were transformed into percent frequency occurrence of a threshall or higher concentration of liquid water content by layer and temperature. That output was then gridded on maps, upon which percent frequency isopleths were drawn. Through the use of accurate liquid water content and temperature climatology, then, a useful tool for predicting the potential for aircraft ising was produced.

THE SMITH-FEDDES LWC MODEL

LWC Model Description. The Smith-Feddes liquid water content model was developed in the mid-1970's at USAFFTAC. It determines liquid water content at a given point in the atmosphere with an empirical model. There are two inputs to the Smith-Feddes model. The first is the 3DNFPH data base, a fifteen-layer vertical cloud distribution data base with a 25 nautical mile (NM) resolution. The 5DNFPH is based on an AFGWC analysis model whose data points contain information on cloud types, bases, tops, percent cloud cover of layers, total cloud cover, present weather, and terrain height. The second input is the Analysis Data Set (ADS), from which ambient temperatures are obtained. This data set has fifteen pressure levels and a coarser 200 NM resolution.

How the LWC Model Works. The Smith-Feddes model extracts moisture data from the first source (3DNEPH) and temperature data from the second source (ADS), creating a liquid water content data base, by layer and temperature, with 200 NM resolution. Liquid water content varies from zero (no cloud) to a maximum of $6~\text{g/m}^3$ (cumulonimbus cloud). Terrain height is taken into account (from 3DNEPH) so that values of each layer are given in meters above ground level (AGL).

PERIOD OF RECORD

Data used for input to the Smith-Feddes model covered a 4-year period of record (1977 to 1980). Because of computer time limitations and the tremendous amount of information contained in the liquid water content data thee, it was impractical to work with more than 4 years of data. Each data point contained about 200 convertions for any 1 month (4 years times 30 days a month times two observations a tay) and over 2,800 discrivations for its annual summary. Therefore, even though only a 4-year period of record was used, the results are statistically representative because of the high observation count.

CHART PREPARATION METHOD

It was necessary to obtain liquid water content data for North America before relating it to a potential light climaticary. The data extracted from the computer runs included point percent frequencies of liquid water context with a 200 NM resolution. These percent frequencies were then potter on a 116,10,200 polar stereographic projection. The data were manually contoured with percent frequency isopictors. Contour intervals were chosen to maintain a similar gradient and concentration of a projector entween product of charts. Manual plotting and contouring allowed climitation of some applicable data prints and also allowed smoothing for easier interpretation.

We also now after the term a descent descent of the particular when the week platted two months, with annual community. These leaves were obtained. In so leaves one countries to 1,524 meters 0.6,000 feets, 1,525 to 1,000 meters 0.0,000 feets, 0.000 meters 0.0,000 feets. All values are relative to a cleve. All values are

Three concentrations of liquid water content were selected as threshold values to simulate three environments for potential icing. If the concentration of liquid water was at or above the threshold value, that observation was counted toward the percent frequency of occurrence. The concentrations used are: .10 grams per cubic meter (g/m^3) (trace to light icing), .50 g/m^3 (moderate icing), and 1.00 g/m^3 (severe icing). In the liquid water content data base, only cumulus and cumulonimbus clouds have concentrations greater than or equal to 1.00 g/m^3 . Therefore, only in these types of clouds would the potential for severe icing be indicated by the liquid water content model. Since most in-cloud severe icing reports occur in cumulus and cumulonimbus clouds, this is desirable.



USING THE CHARTS

The liquid water content charts that follow are climatological data summaries that indicate the potential for aircraft icing. They are intended for use as planning and forecasting aids, not as substitutes for operational icing forecasts, procedures for which are given in AWS/TR-80/001, "Forecasters Guide on Aircraft Icing."

The charts give "percent frequency of occurrence" isopleths at or above a given threshold concentration. Charts are categorized by month, layer, and concentration. The temperature range of all charts is 233 to 273 degrees Kelvin. All layers are above ground level (AGL).

EXAMPLE CHART INTERPRETATION

In Chart 1, the layer shown is surface to 1,524 meters above ground level. The concentration of liquid water content at or below freezing is .10 grams per cubic meter. Since the month is January, winter's colder temperatures and cloudy skies are reflected in the rather high percent frequencies of liquid water content. To interpret the chart, read the isopleths in the same way as a topographic map. New York City is near the 60 percent isopleth. Therefore, New York City's January cloud cover will have a liquid water concentration of .10 grams per cubic meter (or more) about 60 percent of the time. The concentration of .10 grams per cubic meter is not very high and has a high occurrence, but the frequency drops quickly in a moderate (.50 g/m 3) or severe (1.00 g/m 3) icing environment.

The frequencies of liquid water content will be higher than the actual corresponding environments of icing. Even if adequate liquid water content is present at a temperature below freezing, other factors in icing formation (like airspeed or cloud droplet size distribution) may not be conducive to formation of ice on the aircraft.








CHART 1 JANUARY Surface to 1,524 Meters, Concentration $.10~{\rm G/m}^3$




CHART 2 FEBRUARY SURFACE to 1,524 Meters, Concentration $.10 \text{ G/m}^3$




CHART 3 MARCH SURFACE to 1,524 METERS, CONCENTRATION .10 G/M³




CHART 4 APRIL SURFACE TO 1,524 METERS, CONCENTRATION .10 G/M³




CHART 5 MAY SURFACE TO 1,524 METERS, CONCENTRATION .10 G/M³




CHART 6 JUNE SURFACE TO 1,524 METERS, CONCENTRATION .10 G/M³




CHART 7 JULY Surface to 1,524 Meters, Concentration $.10 \text{ G/m}^3$




CHART 8 AUGUST SURFACE to 1,524 Meters, Concentration $.10~{\rm G/m}^3$




CHART 9 SEPTEMBER SURFACE TO 1,524 METERS, CONCENTRATION $,10~\text{G/m}^3$




CHART 10 OCTOBER Surface to 1,524 Meters, Concentration $.10 \text{ G/m}^3$






CHART 11 NOVEMBER SURFACE to 1,524 Meters, Concentration .10 ${\mbox{G/m}}^3$






CHART 12 DECEMBER Surface to 1,524 Meters, Concentration $.10~\text{G/m}^3$




CHART 13 ANNUAL SURFACE TO 1,524 METERS, CONCENTRATION .10 G/M³




CHART 14 JANUARY SURFACE to 1,524 METERS, CONCENTRATION .50 G/M³




CHART 15 FEBRUARY SURFACE TO 1,524 METERS, CONCENTRATION .50 G/M³






CHART 16 MARCH SURFACE TO 1,524 METERS, CONCENTRATION .50 G/M³




CHART 17 APRIL SURFACE TO 1,524 METERS, CONCENTRATION .50 G/M³




CHART 18 MAY SURFACE to 1,524 METERS, CONCENTRATION .50 G/m³




CHART 19 JUNE SURFACE TO 1,524 METERS, CONCENTRATION .50 G/M³




CHART 20 JULY Surface to 1,524 Meters, Concentration .50 g/m³





CHART 21 AUGUST Surface to 1,524 Meters, Concentration $.50 \text{ G/m}^3$



CONTRACT STATES STATES STATES STATES STATES STATES STATES STATES

CHART 22 SEPTEMBER SURFACE to 1,524 Meters, Concentration .50 $\mathrm{G/m^3}$




CHART 23 OCTOBER Surface to 1,524 Meters, Concentration .50 $\mathrm{G/m}^3$




CHART 24 NOVEMBER Surface to 1,524 Meters, Concentration .50 g/m3




CHART 25 DECEMBER Surface to 1,524 Meters, Concentration .50 ${\rm G/m}^3$




CHART 26 ANNUAL SURFACE TO 1,524 METERS, CONCENTRATION .50 G/M³




CHART 27 JANUARY Surface to 1,524 Meters, Concentration $1.00~{\rm G/m}^3$




CHART 28 FEBRUARY SURFACE to 1,524 Meters, Concentration 1.00 g/m³




CHART 29 MARCH Surface to 1,524 Meters, Concentration $1.00~{\rm G/m}^3$




CHART 30 APRIL SURFACE to 1,524 METERS, CONCENTRATION 1.00 G/H³




CHART 31 MAY SURFACE TO 1,524 METERS, CONCENTRATION $1.00~{\rm G/m}^3$




CHART 32 JUNE Surface to 1,524 Meters, Concentration 1.00 g/m³




CHART 33 JULY Surface to 1,524 Meters, Concentration $1.00~{\rm G/m}^3$




CHART 34 AUGUST Surface to 1,524 Meters, Concentration 1.00 ${
m G/m}^3$





CHART 35 SEPTEMBER SURFACE TO 1.524 METERS, CONCENTRATION 1.00 G/M3



<u> 2000.</u>

CHART 36 OCTOBER Surface to 1,524 Meters, Concentration 1.00 ${\rm G/m}^3$




CHART 37 NOVEMBER Surface to 1,524 Meters, Concentration $1.00~{\rm G/m}^3$




CHART 38 DECEMBER Surface to 1,524 Meters, Concentration $1.00~{\rm G/m}^3$




CHART 39 ANNUAL SURFACE TO 1,524 METERS, CONCENTRATION 1.00 G/m³




CHART 40 JANUARY 1,525 to 3,048 Meters, Concentration .10 ${
m G/m}^3$




CHART 41 FEBRUARY 1,525 to 3,048 Meters, Concentration $.10 \text{ G/m}^3$

energias approach somerse exercise because the




CHART 42 MARCH 1,575 to 3,048 Meters, Concentration $.10 \text{ G/m}^3$




CHART 43 APRIL 1,525 to 3,048 Meters, Concentration .10 g/m³




CHART 44 MAY 1,525 to 3,048 Meters, Concentration $.10 \text{ G/m}^3$




CHART 45 JUNE 1,525 to 3,048 Meters, Concentration $.10 \text{ G/m}^3$




CHART 46 JULY 1,525 to 3,048 Meters, Concentration $.10 \text{ G/m}^3$




CHART 47 AUGUST 1,525 to 3,048 Meters, Concentration $.10 \text{ G/m}^3$




CHART 48 SEPTEMBER 1,525 to 3,048 Meters, Concentration $.10~{
m G/M}^3$




CHART 49 OCTOBER 1,525 to 3,048 Meters, Concentration .10 $6/n^3$





CHART 50 NOVEMBER 1,525 to 3,048 Meters, Concentration .10 $\mathrm{G/m}^3$



LONGER SESSEE SPECIES LANGER

CHART 51 DECEMBER 1,525 to 3,048 Meters, Concentration .10 G/m^3





CHART 52 ANNUAL 1,525 to 3,048 Meters, Concentration .10 g/m³



LEGICO COURSERO, PERSONAL CORRESPONDO SUCCESSANDOS SUCCESSANDOS

CHART 53 JANUARY 1,525 to 3,048 Meters, Concentration .50 g/n³





CHART 54 FEBRUARY 1,525 to 3,048 Meters, Concentration .50 g/m³



STATE TOTAL STATE

CHART 55 MARCH 1,525 to 3,048 Meters, Concentration ,50 g/h³




CHART 56 APRIL 1,525 to 3,048 Meters, Concentration $.50 \text{ G/m}^3$




CHART 57 MAY 1,525 to 3,048 Meters, Concentration .50 $\mathrm{G/n}^3$




CHART 58 JUNE 1,525 to 3,048 Meters, Concentration .50 G/m^3




CHART 59 JULY 1,525 to 3,048 Meters, Concentration $.50 \text{ G/m}^3$




CHART 60 AUGUST 1,525 to 3,048 Meters, Concentration .50 $\mathrm{G/m}^3$




CHART 61 SEPTEMBER 1,525 to 3,048 Meters, Concentration .50 $\mathrm{G/m}^{5}$




CHART 62 OCTOBER 1,525 to 3,048 Meters, Concentration .50 $\mathrm{G/m}^3$





CHART 63 NOVEMBER 1,525 to 3,048 Meters, Concentration .50 g/m³

comment recovers between seasons and provides desired length



OWNERS OF MET TO STATE OF THE S






CHART 65 ANNUAL 1,525 to 3,048 Meters, Concentration .50 G/m³





CHART 66 JANUARY 1,525 to 3,048 Meters, Concentration 1.00 $\mathrm{G/m}^3$



WARE CONTRACT STATES

CHART 67 FEBRUARY 1,525 to 3,048 Meters, Concentration 1.00 G/H3





CHART 68 MARCH 1,525 to 3,048 Meters, Concentration $1.00~{\rm G/m}^3$



17**/40** (13) (15/4) (15/4) (15/4) (15/4) (15/4) (15/4) (15/4) (15/4) (15/4) (15/4) (15/4) (15/4) (15/4) (15/4)



STATES ALLEGAS STATES STATES STATES




CHART 70 MAY 1,525 to 3,048 Meters, Concentration $1.00~\mathrm{G/m}^3$




CHART 71 JUNE 1,525 to 3,048 Meters, Concentration $1.00~{\rm G/m}^3$




CHART 72 JULY 1.5% to 3,048 Meters, Concentration 1.00 $\mathrm{G/m^3}$




CHART 73 AUGUST 1,525 to 3,048 Meters, Concentration $1.00~{\rm G/m}^3$




CHART 74 SEPTEMBER 1.525 to 3,048 Meters, Concentration $1.00~{
m G/m}^3$




CHART 75 OCTODER 1,525 to 3,048 Meters, Concentration 1.00 $\mathrm{G/m}^3$




CHART 76 NOVEMBER 1,525 to 3,048 Meters, Concentration 1.00 ${
m G/m}^3$




CHART // DECEMBER : 1325 to 3,048 METERS, CONCENTRATION 1.00 G/N3




CHART 78 ANNUAL 1,525 to 3,048 Meters, Concentration $1.00~\mathrm{G/m}^3$




CHART 79 JANUARY 3,049 to 4,572 Meters, Concentration $.10 \text{ G/m}^3$



STATES STATES STATES STATES STATES




CHART 80 FEBRUARY 3,049 to 4,572 Meters, Concentration $.10~{\rm Gem}^2$




CHART 81 MARCH 3,049 to 4,572 Meters, Concentration $.10 \text{ G/m}^3$





CHART 82 APRIL 3,049 to 4,572 Meters, Concentration .10 G/M³





entities investigate ordinary accepted invitation inter-

CHART 83 MAY 3,049 to 4,572 Meters, Concentration $.10 \text{ G/m}^3$




CHART 84 JUNE 3,049 to 4,572 Meters, Concentration .10 g/m³




CHART % = 30.9 To 4,572 Meters, Concentration .10 s/m³




CHART 86 AUGUST 3,049 to 4,572 Meters, Concentration $.10 \text{ G/m}^3$






CHART 87 SEPTEMBER 3,049 to 4,572 Meters, Concentration $.10 \text{ G/m}^3$





CROCOPY RESOLUTION TEST CHART MATIONAL RUREAU OF STANDARDS 1963 A

BEAREN DOSSESSE SERVED




CHART 88 OCTOBER 3,049 to 4,572 Meters. Concentration ...10 $6/m^3$




CHART 89 NOVEMBER 3,049 to 4,572 Meters, Concentration $.10 \text{ G/m}^3$




CHART 90 DECEMBER 3,049 to 4,572 Meters, Concentration $.10 \text{ G/m}^3$




CHART 91 ANNUAL 3,049 to 4,572 Meters, Concentration $.10 \text{ G/m}^3$




CHART 92 JANUARY 3,049 to 4,572 Meters, Concentration .50 $\mathrm{G/m}^3$




CHART 93 FEBRUARY 3,049 to 4,572 Meters, Concentration .50 g/m³




CHART 94 MARCH 3,049 to 4,572 Meters, Concentration $.50 \text{ G/m}^3$




CHART 95 APRIL 3,049 to 4,572 Meters, Concentration .50 g/h³




CHART 96 MAY 3,049 to 4,572 Meters, Concentration .50 g/m³




CHART 97 JUNE 3,049 to 4,572 Meters, Concentration $.50 \text{ G/m}^3$




CHART 98 JULY 3,049 to 4,572 Meters, Concentration .50 e/h^3




CHART 99 AUGUST 3,049 to 4,572 Meters, Concentration .50 G/h^3




CHART 100 SEPTEMBER 3,049 to 4,572 Meters, Concentration .50 G/m^3




CHART 101 OCTOBER 3,049 to 4,572 Meters, Concentration .50 e/h^3

CHICAGO BOOCAGO DISTINO





CHART 102 NOVEMBER 3,049 to 4,572 Meters, Concentration .50 g/m 3



E

CHART 103 DECEMBER 3,049 to 4,572 Meters, Concentration ,50 g/m³




CHART 104 ANNUAL 3,049 to 4,572 Meters, Concentration .50 G/m^3




CHART 105 JANUARY 3,049 to 4,572 Meters, Concentration 1.00 g/m³





CHART 106 FEBRUARY 3,049 to 4,572 Meters, Concentration $1.00~\text{G/m}^3$



CONTRACTOR CONTRACTOR CONTRACTOR CONTRACTOR CONTRACTOR

CHART 107 MARCH 3,049 to 4,572 Meters, Concentration 1.00 g/m³




CHART 108 APRIL 3,049 to 4,572 Meters, Concentration 1.00 G/H3




CHART 109 MAY 3,049 to 4,572 Meters, Concentration 1.00 $\mathrm{g/m}^3$




CHART 110 JUNE 3,049 to 4,572 Meters, Concentration 1.00 G/H³




CHART 111 JULY 3,049 to 4,572 Meters, Concentration 1.00 $\mathrm{G/m}^3$




CHART 112 AUGUST 3,049 to 4,572 Meters, Concentration $1.00~{\rm G/m}^3$




CHART 113 SEPTEMBER 3,049 to 4,572 Meters, Concentration 1.00 g/m 3




CHART 114 OCTOBER 3,049 to 4,572 Meters, Concentration $1.00~{\rm G/M}^3$




CHART 115 NOVEMBER 3,049 to 4,572 Meters, Concentration $1.00~{\rm G/M}^3$




CHART 116 DECEMBER 3,049 to 4,572 Meters, Concentration $1.00~{\rm G/m}^3$




CHART 117 ANNUAL 3,049 to 4,572 Meters, Concentration 1.00 g/m³

BIBLIOGRAPHY

- Feddes, Robert G., Development of a Gridded Data Base, USAFETAC IN 74-2, USAF Environmental Technical Applications Center, Scott Air Force Base, Illinois, 62226, April 1974.
- Feddes, Robert G., A Synoptic-Scale Model for Simulating Condensed Atmospheric Moisture, USAFETAC TN 74-4, USAF Environmental Technical Applications Center, Scott Air Force Base, Illinois, 62225, June 1974.
- Forecasters' Juide on Aircraft Icing, AWS/TR-80/001, Air Weather Service, Scott Air Force Base, Illinois, 62225, March 1980.
- Fye, Falko K., The AFGWC Automated Cloud Analysis Model, AFGWC TM 78-002, Air Force Global Weather Central, Offutt Air Force Base, Nebraska, 68113, June 1978.
- Jackson, G.C., Icing Climatology for Northern Europe, AFWAL-TM-80-16-FSW, Air Force Wright Aeronautical Laboratories, Wright-Patterson Air Force Base, Ohio, 45433, February 1980.
- Smith, Robert D., <u>Atmospheric Moisture Parameterization</u>, USAFETAC TN 74-1, USAF Environmental Technical Applications Center, Scott Air Force Base, IllInois, 62225, January 1974.



DISTRIBUTION

The second secon
AWS/DN, Scott AFB, IL 62225-50082
AWS/DO, Scott AFB, IL 62225-5008
AWS/SY, Scott AFB, IL 62225-5008
OL A. HQ AWS, Buckley ANG Base, Aurora, CO 80011-9599
OL-C, HQ AWS, Chanute AFB, IL 61868-5000
OL-E, HQ AWS, ATZL-CAW-E, Ft Leavenworth, KS 66027-5300
On the no new door the term of the Least of
SD/YDA, PO Box 92960, Worldway Postal Ctr, Los Angeles, CA 90009-2960
OL-G, HQ AWS, NHC Rm 631, 1320 S Dixie Hwy, Coral Gables, FL 33146
OL-H, HQ AWS (ATSI-CD-CS-SWO), Ft Huachuca, AZ 85613-5000
OL-I, HQ AWS, Ft Monroe AIN, VA 23651-5000
AFOTEK OL-NX, 1313 Halley Circle, Norman, OK 73069
OL-L, HQ AWS, Keesler AFB, MS 39534-5000
SM-ALC/MMA, McClellan AFB, CA 95652-5000.
Det 1, HQ AWS, Pentagon, Washington, DC 20330-6560
Det 2, HQ AWS, Pentagon, Washington, DC 20330-5054
AFSCF/WE, PO Box 3430, Sunnyvale AFS, CA 94088-3430
Det 6, HQ AWS, Langley AFB, VA 23665-500040
Det 8, HQ AWS, PO Box 4239N, Las Vegas, NV 890301
Det 9, HQ AWS, PO Box 12297, Las Vegas, NV 89112-02971
1WW/DN, Hickam AFB, HI 96853-5000
20WS/DN, APO San Francisco 96328-5000
30WS/DN, APO San Francisco 96301-0420
2WW/DN, APO New York 09012-5000
7WS/DN, APO New York 09403-500023
28WS/DN, APO New York 09127-500010
31WS/DN, APO New York 09223-5000
3WW/DN, Offutt AFB, NE 68113~50001
9WS/DN, March AFB, CA 92518-5000
11WS/DN, Elmendorf AFB, AK 99506-5000
24WS/DN, Randolph AFB, TX 78150-5000
26WS/DN, Barksdale AFB, LA 71110-5002
4WW/DN, Peterson AFB, CO 80914-5000.
2WS/DN, Andrews AFB, MD 20334-5000
5WW/DN, Langley AFB, VA 23665-5000
1WS, MacDill AFB, FL 33608-5000
3WS/DN, Shaw AFB, SC 29152-5000
5WS/DN, Ft McPherson, GA 30330-5000
25WS/DN, Bergstrom AFB, TX 78743-5000
AFGWC/SDSL, Offutt AFB, NE 68113-5000
7WW/DN, Scott AFB, IL 62225-5008
6WS, Tinker AFB, OK 73145-5000
15WS/DN, McGuire AFB, NJ 08641-5002
17WS/DN, Travis AFB, CA 94535-500014
3350 TECH TG/TTGU-W, Stop 62, Chanute AFB, IL 61868-5000
Naval Research Laboratory, Code 4323, Washington, DC 20375
NAVOCEANCOMFAC, NSTL, Bay St Louis, MS 39529-5002
COMNAVOCEAN, NSTL, MS 39529-50002
NAVOCEANO, ATTN: Code 9220 (Tony Ortolano), Bay St Louis, MS, NSTL 39522-50012
NEPRF, Monterey, CA 93943-5106
Naval Post-Graduate School, NC4(63rd), Monterey, CA 93943-5100
Naval Western Oceanography Ctr, Box 113, Attn: Tech Library, Pearl Harbor, HI 96860-50001
Naval Oceanography Command Ctr, COMNAVMAR Box 12, Attn: AG1 Karen Seib,
FPO San Francisco, CA 96630-5000
-1/5 Can 1/2004 - 20 //030 200444444444444444444444444444444444
AFGI Library Attn. SULLR Stop 20 Hangoom AFR MA 01721-5000
AFGL Library, Attn: SULLR, Stop 29, Hanscom AFB, MA 01731-5000
AFGL Library, Attn: SULLR, Stop 29, Hanscom AFB, MA 01731-5000
AFGL Library, Attn: SULLR, Stop 29, Hanscom AFB, MA 01731-5000
AFGL Library, Attn: SULLR, Stop 29, Hanscom AFB, MA 01731-5000
AFGL Library, Attn: SULLR, Stop 29, Hanscom AFB, MA 01731-5000
AFGL Library, Attn: SULLR, Stop 29, Hanscom AFB, MA 01731-5000

⇒ U.S. GOVERNMENT PRINTING OFFICE : 1986—652-005/24117

12-86

07/